

A Case Report on Chikungunya with Diagnostic Evaluation & Management

**Ms. Alfiya shaikh¹, Ms. Pallavi Dhole², Aniket Pathade³, P.S. Pande⁴
, Mrs. Kavita Gomase⁵**

1] Ms. Alfiya shaikh, G.N.M 2nd year, Florence Nightingale Training College of Nursing, Datta Meghe Institute of Medical Science (DU) Sawangi (M) Wardha. Email: -alfishaikh701@gmail.com, Mobile no: 8767271807

2] Ms. Pallavi Dhole, Nursing Tutor, Florence Nightingale Training College of Nursing, Datta Meghe Institute of Medical Science (DU) Sawangi (M) Wardha. Email: pallavidhole2007@gmail.com, Mobile no: 9960597125

3] Research Scientist, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (M), Wardha, Maharashtra, India.

4] Dept. of Chemistry, Yeshwantrao Chavan College of Engineering, Nagpur
pournimapande@yahoo.co.in

5] Assistant Professor, Dept of Obstetrics and Gynecology Nursing, Smt. Radhikabai Meghe Memorial College of Nursing, kavitapraavin3@gmail.com

Corresponding Author : Ms. Alfiya shaikh

Ms. Alfiya shaikh, G.N.M 2nd year, Florence Nightingale Training College of Nursing, Datta Meghe Institute of Medical Science (DU) Sawangi (M) Wardha.

ABSTRACT

Introduction: Infection with the Chikungunya virus is spread via mosquitoes. The chikungunya virus is to blame (CHIKV). Fever and arthralgia are the most common symptoms. Chikungunya virus is spread by *Aedes aegypti* and *Aedes albopictus* mosquitoes, which are found throughout the Asia, Europe, Africa, and America are the four continents. The same mosquitoes that cause Chikungunya also spread Dengue and Zika viruses, and there have been occurrences of coinfections. To tackle the infection, there is currently no vaccine or specific therapy available. The goal of treatment is to reduce the severity of the disease's symptoms. **Patient Information:** A 33-year-old male was admitted to AVBR hospital with a chief complaint of fever, & severe joint pain for 3 days, headache, muscle pain, nausea & vomiting. **Main symptoms & important clinical finding:** -He has the chief complaint of fever, & severe joint pain for 3 days, headache, muscle pain, Nausea & vomiting after careful history collection, physical examination, Complete blood count, RBS, CT scan, MRI Etc. my patient diagnoses as chikungunya. **Medical Management:** -A patient provides a medicine as per doctor's order like antipyretics, analgesics such as Tab. Paracetamol, Tab. acetaminophen, Tab. Emset. **Nursing Perspectives:** - Monitoring vital signs hourly. Prescribe medicine as per doctor's order. maintain IV fluid 4 hourly & record intake output chart. maintain records & reports. **Conclusion:** Early Diagnosis & Timely Treatment patient condition was improved.

Keywords: -Chikungunya virus infection, *Aedes aegypti*, Dengue.

Introduction

The chikungunya virus, a type of alphavirus, causes chikungunya fever, a mosquito-borne sickness. *Aedes aegypti* and *Aedes albopictus* mosquitoes are the predominant chikungunya vectors throughout Asia and the Indian Ocean islands. In 1952, the virus was first detected in Tanzania. It's been linked to a series of outbreaks in several nations since then. The virus has been detected in Africa, Southeast Asia, and India, among other places. Several countries in the affected regions are reporting sporadic instances regularly. Chikungunya is now found in over a hundred countries throughout the world.[1]

Chikungunya is a mosquito-borne virus that was first detected in 1952 in southern Tanzania after an outbreak. It's an RNA virus that belongs to the alphavirus genus of the *Togaviridae* family.[2]

The term "chikungunya" is derived from a Kimakonde expression that means "to become distorted," & alludes to joint pain sufferers' bent posture (arthralgia). Chikungunya fever is a febrile illness that can afflict anyone of any age and is usually self-limiting. Epidemic peaks tend to diminish as population immunity improves because CHIKV infection gives long-term protection.[3]

Chikungunya is a virus spread by arthropods that causes severe polyarthralgia and arthritis in people. It is found in Africa, Southeast Asia, and India. Six Bangladeshi individuals with chikungunya fever are discussed in this briefcase series. Although Bangladesh is in an endemic zone, the disease is uncommon here, necessitating prompt diagnosis and treatment.[4]

In late 2013, in the Caribbean islands, the Chikungunya virus was detected for the first time in the Americas. Tourists who have been infected can transfer the disease to other locations. The chikungunya virus has no vaccine or medication available to prevent or treat the infection. Mosquito bite prevention is a good approach for travelers to protect themselves against mosquito bites. When visiting countries where the chikungunya virus is present, use insect repellent, wear long sleeves and trousers, and stay in areas with air conditioning or window and door screens.[5]

The rash appears in 40–50% of cases, usually as a maculopapular rash two to five days following the onset of symptoms. Abdominal pain, nausea, vomiting, or diarrhea are all possible digestive symptoms. Significant weariness and pain hinder normal activity in more than half of cases. Eye inflammation in the form of iridocyclitis or uveitis, as well as retinal abnormalities, occurs seldom. There's a chance you'll get liver damage for a while.[6]

Patient information

A 33-year-old male admitted to AVBR hospital with a chief complaint of fever, & severe joint pain for 3 days, headache, muscle pain, nausea & vomiting. He was diagnosed as Chikungunya after undergoing all investigations such as complete blood count, CT scan, MRI.

The primary concern and symptoms of the patient

A chief complaint of fever, & severe joint pain for 3 days, headache, muscle pain, nausea & vomiting. He was diagnosed as Chikungunya after undergoing all investigations such as complete blood count, CT scan, MRI.

Medical family and psycho-social history

The patient suffering from chikungunya for 3 days in family history he belongs to a nuclear family. His monthly family income is 50,000. In this family, there are no medical problems like Hypertension, Diabetes mellitus, asthma. There are no hereditary or genetic disorders in the family. He was mentally stable. He is oriented to date, time and place and he maintains a good relationship with family members.

Clinical Finding

On clinical examination, the patient has high fevers, discoloration in the eye, conjunctivitis is all found. In rare circumstances, cervical or widespread lymphadenopathy has been described. Severe arthralgia, myalgia, and rash are other common signs.[7]

Timeline

A 33-year-old male was admitted to AVBR hospital with a chief complaint of fever, & severe joint pain for 3 days, headache, muscle pain, nausea & vomiting. Early diagnosis & proper treatment of my patient's condition was improved.

Diagnosis Assessment

Based on careful history collection, physical examination and investigation, a patient having high-grade fever, shortness of breathing is present. My patient's all investigation done like complete blood count, Routine test, random blood sugar, renal function test, hemogram, urine routine, ultrasound, CT scan, MRI (if needed) were all performed. Serological techniques, like enzyme-linked immunosorbent assays (ELISA), can be used to confirm the presence of anti-chikungunya antibodies, including IgM and IgG.

Diagnostic challenging

No challenging during diagnosis evaluation. After history collection and physical examination and all investigations a doctor's diagnosis is a case of chikungunya.

Therapeutic Investigation:

In chikungunya virus infection, there is no specific treatment. Options include rest, beverages, and the use of nonsteroidal anti-inflammatory drugs (NSAIDs) to relieve acute pain and fever. NSAIDs, corticosteroids, and physiotherapy may be used to alleviate persistent joint pain. During the first week of sickness, patients infected with chikungunya should be protected from future mosquito exposure to reduce the risk of local transmission. He was looking at all treatments and the outcome was good. His signs and symptoms were reducing. No change in therapeutic interventions.

Nursing Perspectives

Monitoring vital signs hourly & recorded. Prescribe medicine as per doctors' orders. Maintain IV fluid 4 hourly & record intake output chart, monitor vital signs, providing patient and relatives with psychological support, Establishing good interpersonal relationship. Maintaining confidentiality will relieve the anxiety of the patient.

and relatives. It will help in avoiding conflict with physicians, nurses and other health care providers. Drinks plenty of water to avoid dehydration. Reduce fever and pain with paracetamol or acetaminophen.

Follow up and outcomes

The patient was planned for follow-up regularly on basis of advice given by the physician.

Preventive Measures

Putting on long-sleeved shirts, long pants, and other skin-protective clothing. Applying insect repellent to the skin or clothing. Having enough screens in indoor rooms to keep mosquitoes at bay. If you sleep during the day, cover your bed with insecticide-treated mosquito nets. Cover your face and neck with mosquito netting, as well as using gloves or repellents. Avoid travel to areas where the chikungunya virus is circulating. Using mosquito coils and insecticide vaporizers during the daytime. Garbage should be stored in the proper place or in plastic bags. Getting rid of old tires collect or place left outside. Draining water from a container such as potted plants, buckets, rain gutters.

Discussion

The authors present a case of rhabdomyolysis caused by Chikungunya virus infection, a rather uncommon infectious consequence. With conservative management, the patient healed without incident and was discharged to his home. [8]

Temperature (92 percent), which can range from mild to severe & arthralgia 87%, are the most common presenting characteristics recorded in the literature. The discomfort in the joints is usually worse first thing in the morning. Mild exercise relieves it, but vigorous movements increase it. In 62 percent of cases, patients have headaches and chills. Chikungunya fever can be diagnosed with a Chikungunya virus serology or real-time reverse transcription-polymerase chain reaction (RT-PCR). [9]

Acute joint pain is the most prevalent symptom of chikungunya fever, and it occurs in nearly every clinical episode. We saw the same thing in both of our circumstances. The arthralgia is frequently symmetrical and peripheral, affecting both minor and large joints in the hands. [10]

As a result of the severe discomfort and swelling in the joints, patients generally experience pain that lasts for weeks to months and is incapacitating. Although certain Arthralgia caused by chikungunya fever could last for years. The majority of illnesses endure only a few weeks or months. Only two people in our case study had ongoing joint discomfort two to three months after the illness had cleared up. [11,17]

Many tropical and subtropical locations around the world, including Sub-Saharan Africa, Southeast Asia, India, and the western Pacific, are endemic for the virus. Throughout recent years, Chikungunya has become widespread in the Caribbean, with a peak in 2014, when many cases were documented in Haiti, the Dominican Republic, and other parts of the region. [12,18,19]

Vector-borne transmission to humans is the most common mode of transmission. There is yet to be any proof of transfer from person to person. The CHIK virus, on the other hand, can be aerosolized, and laboratory workers have been infected as a result of working with it. [13]

The severe nature of CHIKV rheumatic disease, which can have a substantial influence on affected patients' quality of life for weeks, months, or even years. The recent CHIKV outbreaks, as well as their shown potential to move fast into new areas, have reignited interest in developing measures to prevent or treat CHIKV-related disease. [14,15,16]

Conclusion

A 33-year-old was admitted to AVBR hospital with a chief complaint of fever, & severe joint pain in the last 3 days, headache, muscle pain, nausea & vomiting. The patient had undergone all investigations and after that, she was diagnosed with chikungunya.

Reference

1. Weaver SC, Lecuit M. Chikungunya virus and the global spread of a mosquito-borne disease. *New England Journal of Medicine*. 2015 Mar 26;372(13):1231-9.
2. Uddin KS. Chikungunya-yet another mosquito borne epidemic burden for Bangladesh. *KYAMC Journal*. 2017 Aug 31;8(1):1-3.
3. Marques CD, Ranzolin A, Cavalcanti NG, Duarte AL. Arboviruses related with chronic musculoskeletal symptoms. *Best Practice & Research Clinical Rheumatology*. 2020 Aug 1;34(4):101502.
4. Hassan R, Rahman MM, Moniruzzaman M, Rahim A, Barua S, Biswas R, Biswas P, Mowla SG, Chowdhury MJ. Chikungunya—an emerging infection in Bangladesh: a case series. *Journal of medical case reports*. 2014 Dec;8(1):1-3.

5. Pastula DM, Smith DE, Beckham JD, Tyler KL. Four emerging arboviral diseases in North America: Jamestown Canyon, Powassan, chikungunya, and Zika virus diseases. *Journal of neurovirology*. 2016 Jun;22(3):257-60.
6. Lierl M. Periodic fever syndromes: a diagnostic challenge for the allergist. *Allergy*. 2007 Dec;62(12):1349-58.
7. Wallace MR, Hale BR, Utz GC, Olson PE, Earhart KC, Thornton SA, Hyams KC. Endemic infectious diseases of Afghanistan. *Clinical Infectious Diseases*. 2002 Jun 15;34(Supplement_5): S171-207.
8. Elfert KA, Abdelwahed M, Chi G. Chikungunya virus infection-related rhabdomyolysis: a case report. *Cureus*. 2019 Feb;11(2).
9. Elfert KA, Abdelwahed M, Chi G. Chikungunya virus infection-related rhabdomyolysis: a case report. *Cureus*. 2019 Feb;11(2).
10. Hassan R, Rahman MM, Moniruzzaman M, Rahim A, Barua S, Biswas R, Biswas P, Mowla SG, Chowdhury MJ. Chikungunya—an emerging infection in Bangladesh: a case series. *Journal of medical case reports*. 2014 Dec;8(1):1-3.
11. Hassan R, Rahman MM, Moniruzzaman M, Rahim A, Barua S, Biswas R, Biswas P, Mowla SG, Chowdhury MJ. Chikungunya—an emerging infection in Bangladesh: a case series. *Journal of medical case reports*. 2014 Dec;8(1):1-3.
12. Weaver SC, Forrester NL. Chikungunya: Evolutionary history and recent epidemic spread. *Antiviral research*. 2015 Aug 1; 120:32-9.
13. Gamage SD, Kravolic SM, Roselle G. Emerging infectious diseases: concepts in preparing for and responding to the next microbial threat. *Koenig and Schultz's disaster medicine: Comprehensive principles and practices*. 2009:75-102.
14. Morrison CR, Plante KS, Heise MT. Chikungunya virus: current perspectives on a reemerging virus. *Microbiology spectrum*. 2016 May 13;4(3):4-3.
15. Abbafati, Cristiana, Kaja M. Abbas, Mohammad Abbasi, MitraAbbasifard, Mohsen Abbasi-Kangevari, HedayatAbbastabar, FoadAbd-Allah, et al. “Five Insights from the Global Burden of Disease Study 2019.” *LANCET* 396, no. 10258 (October 17, 2020): 1135–59.
16. Abbafati, Cristiana, Kaja M. Abbas, Mohammad Abbasi, MitraAbbasifard, Mohsen Abbasi-Kangevari, HedayatAbbastabar, FoadAbd-Allah, et al. “Global Burden of 369 Diseases and Injuries in 204 Countries and Territories, 1990-2019: A Systematic Analysis for the Global Burden of Disease Study 2019.” *LANCET* 396, no. 10258 (October 17, 2020): 1204–22.
17. Franklin, Richard Charles, Amy E. Peden, Erin B. Hamilton, Catherine Bisignano, Chris D. Castle, Zachary Dingels V, Simon Hay I, et al. “The Burden of Unintentional Drowning: Global, Regional and National Estimates of Mortality from the Global Burden of Disease 2017 Study.” *INJURY PREVENTION* 26, no. SUPP_1, 1 (October 2020): 83–95. <https://doi.org/10.1136/injuryprev-2019-043484>.
18. James, Spencer L., Chris D. Castle, Zachary Dingels V, Jack T. Fox, Erin B. Hamilton, Zichen Liu, Nicholas L. S. Roberts, et al. “Estimating Global Injuries Morbidity and Mortality: Methods and Data Used in the Global Burden of Disease 2017 Study.” *INJURY PREVENTION* 26, no. SUPP_1, 1 (October 2020): 125–53. <https://doi.org/10.1136/injuryprev-2019-043531>.
19. James, Spencer L., Chris D. Castle, Zachary Dingels V, Jack T. Fox, Erin B. Hamilton, Zichen Liu, Nicholas L. S. Roberts, et al. “Global Injury Morbidity and Mortality from 1990 to 2017: Results from the Global Burden of Disease Study 2017.” *INJURY PREVENTION* 26, no. SUPP_1, 1 (October 2020): 96–114. <https://doi.org/10.1136/injuryprev-2019-043494>.